

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Journal of Experimental Marine Biology and Ecology

journal homepage: www.elsevier.com/locate/jembe

UVR-induced photosynthetic inhibition dominates over DNA damage in marine dinoflagellates exposed to fluctuating solar radiation regimes

E. Walter Helbling^{a,b,*}, Anita G.J. Buma^c, Willem van de Poll^c,
M. Verónica Fernández Zenoff^{b,d}, Virginia E. Villafañe^{a,b}^a Estación de Fotobiología Playa Unión, Casilla de Correos N° 15 (9103) Rawson, Chubut, Argentina^b Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)^c Department of Marine Biology – Centre for Ecological and Evolutionary Studies, University of Groningen, P.O. Box 14, 9750 AA Haren, The Netherlands^d Planta Piloto de Procesos Industriales Microbiológicos (PROIMI-CONICET), Avda. Belgrano y Pje. Caseros (4000) San Miguel de Tucumán, Argentina

ARTICLE INFO

Article history:

Received 18 June 2008

Received in revised form 30 July 2008

Accepted 30 July 2008

Keywords:

Dinoflagellates

DNA

Mixing

Patagonia

photosynthesis

UVR

ABSTRACT

The combined effect of solar radiation (UV-B (280–315 nm), UV-A (315–400 nm) and PAR (400–700 nm)) and vertical mixing (i.e., fluctuating radiation regimes) on the marine dinoflagellates *Gymnodinium chlorophorum*, *Heterocapsa triquetra* and *Prorocentrum micans* was investigated during the austral spring in Patagonia, Argentina. Photosynthesis, measured as radiocarbon incorporation, and accumulation of DNA damage, as cyclobutane pyrimidine dimers (CPDs), were investigated under simulated mixed and non-mixed water column conditions using 3 h incubations centered at local noon. Static samples had significant UVR-induced photoinhibition that was higher in *H. triquetra* as compared to the other two species. Increasing mixing speed significantly increased UVR-induced inhibition of carbon fixation in *G. chlorophorum* and *H. triquetra*. No significant UVR effect was observed in *P. micans* under any of the mixing regimes. Most of the loss in carbon fixation in *G. chlorophorum* was due to UV-B while in *H. triquetra* it was due to UV-A. Part of these responses may be associated to the presence of UV-absorbing compounds which were abundant in *P. micans*, and low in *H. triquetra* and in *G. chlorophorum*. However, other variables such as cell size and active repair might have also influenced our results. We did not detect CPD accumulation in any of the species, probably because of the low solar angle that resulted in very low levels of DNA effective UV-B dose. Our results indicate that exposure to solar UVR in the Patagonia area during spring time (even during ozone depletion events) has a clear impact on photosynthesis and much less or negligible on DNA in the three studied species.

© 2008 Elsevier B.V. All rights reserved.

* Corresponding author.

E-mail address: whelbling@efpu.org.ar (E.W. Helbling).